

Igor V. Bayak

A lemma on the minimal surfaces

Stated lemma contains the assertions about isomorphism of exact m-forms and exterior differentials of regular m-maps, of linearly harmonic m-forms and exterior differentials of regular harmonic m-maps, of global minimal $(n-m)$ -surfaces and level $(n-m)$ -surfaces of regular minimal m-maps.

Let $I = \{1, \dots, n\}$; $J = \{1, \dots, m\}$, $1 \leq m < n$; $(J) \equiv (i_1, \dots, i_j, \dots, i_m)$: $J \rightarrow I$ - is an arbitrary ranked in ascending order arrangement, i.e. such injection that $i_{j-1} < i_j$; $\{(J)\}$ - is set of all ordered injections; E - is n -dimensional Euclidean space; $(\bar{e}^J)_I$ - is basis of E ; $(\hat{e}^{(J)})_{\{(J)\}} \equiv (dx_{i_1} \wedge \dots \wedge dx_{i_j} \wedge \dots \wedge dx_{i_m})_{\{(J)\}}$ - is basis of $\Lambda^m E^*$;

$$\varphi^m : E \rightarrow R^m \left| \lim_{\sup \{|\Delta x_j|_J\} \rightarrow 0} \left| \varphi_i^m(\bar{x} + \Delta \bar{x}) - \varphi_i^m(\bar{x}) \right| \rightarrow 0 \right.$$

$$\left. \lim_{\sup \{|\Delta x_j|_J\} \rightarrow 0} \frac{\varphi_i^m(\bar{x}) - \varphi_i^m(\bar{x} + \Delta \bar{x})}{\varphi_i^m(\bar{x}) - \varphi_i^m(\bar{x} - \Delta \bar{x})} \rightarrow 1 \quad (\forall \bar{x} \in E, i \in J) \right.$$

is an arbitrary m-map, where simultaneously are satisfied the conditions of continuity

and existence of differential. If $J(\bar{x}) \equiv \left(\frac{\partial \varphi_i^m(\bar{x})}{\partial x_j} \right)_{i \in J, j \in I}$ - is Jacobi matrix of the map

φ^m in the point $\bar{x}$ and $\text{rank } J(\bar{x}) = m \quad \forall \bar{x} \in E$ (that is equivalent to the existence of m linear independent gradients in all Euclidean space), then m-map is called by regular in E . Let $\{\varphi^m\}$ - is space of all regular m-maps, then $\varphi_{(n-m)}(\bar{x}_0) \equiv \{\bar{x} \mid \varphi^m(\bar{x}) = \varphi^m(\bar{x}_0)\}$ - is level $(n-m)$ -surface corresponding to the regular map φ^m and to the point of belonging $\bar{x}_0$; $\varphi_{(n-m)} \equiv \{\varphi_{(n-m)}(\bar{x}_0)\}_{\bar{x}_0 \in E}$ - is $(n-m)$ -foliation of E and $\{\varphi_{(n-m)}\}$ - is space of $(n-m)$ -foliations of E . In addition

we define that $d\varphi^m(\bar{x}) \equiv \sum_{\{(J)\}} \det \left(\frac{\partial \varphi_i^m(\bar{x})}{\partial x_j} \right)_{j \in (J)}^{i \in J} \hat{e}^{(J)}$ - is exterior differential of φ^m in

the point $\bar{x}$; $d\varphi^m \equiv \{d\varphi^m(\bar{x})\}_{\bar{x} \in E}$ - is exterior differential of φ^m in E ; $\{d\varphi^m\}$ - is

space of exterior differentials in E . Let $H^i(\bar{x}) \equiv \left(\frac{\partial^2 \varphi_i^m(\bar{x})}{\partial x_j \partial x_k} \right)_{j, k \in I}$, where $i \in J$, - are

Hesse matrices of φ^m in the point $\bar{x}$, then $\{\eta^m\} \equiv \{\varphi^m \mid \text{Sp} H^i = 0 \quad (\forall i \in J)\}$ - is space

of all regular harmonic m-maps. Let also $H_{(J)}^i(\bar{x}) \equiv \left(\frac{\partial^2 \varphi_i^m(\bar{x})}{\partial x_j \partial x_k} \right)_{j, k \in I \setminus (J)}$, where

$i \in J, (J) \in \{(J)\}$, - are sectional Hesse matrices of φ^m in the point $\bar{x}$, then $\{\mu^m\} \equiv \{\varphi^m \mid \text{Sp} H_{(J)}^i = 0 \quad (\forall i \in J, (J) \in \{(J)\})\}$ - is space of all regular minimal m-maps, in

particular, $\{\mu^1\} \equiv \{\varphi^1 \mid \text{Sp} H_i^1 = 0 (\forall i \in I)\}$ - is space of all regular minimal 1-maps. Further let $\{\hat{\varphi}^m\}$ - is space of all exact m-forms and $\{\hat{\eta}^m\}$ - is space of all linearly harmonic m-forms, i.e. such m-forms that vanish by linear operator $(d + \delta)$, and then hold the affirmation.

Lemma:

1. $\{d\varphi^m\} \approx \{\hat{\varphi}^m\}$
2. $\{d\eta^m\} \approx \{\hat{\eta}^m\}$
3. Space of foliations $\{\mu_{(n-m)}\}$ is isomorphic to space of foliations of E by global minimal $(n-m)$ -surfaces, i.e. the minimality condition of regular m-map serves as the differential criterion of extremality (minimality) of integral functional of volume of surfaces under local variations.